



Original Full Paper

Mycoplasma bovis pneumonia in feedlot cattle and dairy calves in Argentina

Carlos A. Margineda^{1,2*}, Gustavo O. Zielinski^{1,2}, Susana Jurado³, Ferrella Alejandra⁴,
 Marina Mozgovej⁴, Ana C. Alcaraz⁵, Alfonso López⁶

¹Grupo de Sanidad Animal, Estación Experimental Agropecuaria INTA Marcos Juárez, Córdoba, Argentina.

²Facultad de Ciencias Veterinarias, Universidad Nacional de Rosario, Casilda, Santa Fe, Argentina.

³Servicio Central de Microscopía Electrónica, Facultad de Ciencias Veterinarias, Universidad Nacional de La Plata, La Plata, Buenos Aires, Argentina.

⁴Instituto de Virología, CICVyA, INTA Castelar, Morón, Buenos Aires, Argentina.

⁵College of Veterinary Medicine, Western University of Health Sciences, Pomona, California, USA.

⁶Department of Pathology and Microbiology, Atlantic Veterinary College, University of Prince Edward Island, Charlottetown, Prince Edward Island, Canada.

*Corresponding author: National Institute of Agriculture Technology, Córdoba, Argentina E-mail: margineda.carlos@inta.gov.ar

Submitted February 13th 2017, Accepted June 3rd 2017

Abstract

Mycoplasma bovis has emerged as an important cause of feedlot pneumonia in many countries. The aim of this paper is to describe six cases of bovine *Mycoplasma* pneumonia in five different premises in Argentina. Gross examination revealed chronic bronchopneumonia with multiple foci of caseous necrosis. Microscopically, these contained a necrotic center with abundant hyper eosinophilic granular material surrounded by granulation tissue. Affected lung tested positive for *M. bovis* by immunohistochemistry and electron microscopy revealed membranous structures compatible with *Mycoplasma* spp. To our knowledge, this is the first report of *M. bovis* pneumonia in Argentina.

Key words: chronic bronchopneumonia, electron microscopy, feedlot, immunohistochemistry, *Mycoplasma bovis*.

Introduction

Bovine respiratory disease (BRD) is the most important cause of economic loss of the beef industry in many parts of the world (12, 14, 26). It is a multifactorial condition that involves host and environmental factors, and several viruses and bacteria. The most common viruses associated in cattle with BRD are bovine herpesvirus (BHV-1), bovine parainfluenza virus 3 (BPIV-3), bovine viral diarrhea virus 1 and 2 (BVDV-1/2), bovine respiratory syncytial virus (BRSV) and bovine coronavirus (BCoV) (5, 14, 24). These viral infections are usually transient but predispose to opportunistic secondary bacterial pneumonia (8, 22, 24). *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni*, and *Mycoplasma bovis* are the most common bacterial isolates

recovered from pneumonic lungs in outbreaks of BRD worldwide (2, 14, 15, 24, 25).

For the last 15 years, *M. bovis* has emerged as one of the most important bacterial pathogens involved in BRD worldwide (9, 23, 26, 29). Although this bacterium is a commensal nasopharyngeal organism in healthy cattle, under certain conditions, it can cause pneumonia, arthritis, synovitis, keratoconjunctivitis, mastitis, metritis, abortion, infertility, otitis and skin abscesses (8, 23). In the lung, *M. bovis* produces chronic bronchopneumonia characterized by caseonecrotic nodules in the cranioventral lung lobes (2, 10, 24). Microscopic findings suggest that lung necrosis is originated in small bronchi and bronchioles (15, 18). The persistence of the agent in the lung tissue and the chronicity of the pulmonary lesions imply that the immune response is ineffective in eliminating the infection (6, 18, 21).

Epidemiological studies indicate that the incidence of *M. bovis* pneumonia has notably increased in many countries such as the United States of America, Canada, Mexico, France, Northern Ireland and Italy, particularly in feedlots. (2, 5, 14, 19, 23, 25, 26, 30). In Argentina, *M. bovis* has never been previously described as a cause of BRD outbreaks or pneumonia.

The objective of this report is to describe six outbreaks of *M. bovis* pneumonia in feedlot steers and dairy calves in four different herds in the Argentinean pampas.

Materials and methods

The first outbreak of BRD associated with *M. bovis* in Argentina occurred in dairy calves at a farm with 220 milking cows in May 2012 (Farm A). In 2013, five other outbreaks took place in three different industrial feedlots (Farms B, C, and D) where animals had arrived for fattening from the north and central Provinces of Argentina. Other relevant information collected from the farms and veterinarians included fatal disease onset (FDO), defined as the period between feedlot arrival and first treatment for pneumonia; the type and frequency of antibiotic treatment; and the time interval between feedlot arrival and the day of death (DD).

Three calves that were unresponsive to antimicrobial treatment for pneumonia (calves 1, 4, and 5) were euthanized (Fig. 1), while three others (calves 2, 3, and 6) were found dead. All animals were necropsied, and lung samples were collected for bacteriological culture, virus isolation, histopathology and immunohistochemistry (IHC), and electron microscopy.

The bacteriological analysis was done in duplicates, inoculating lung tissue on blood agar and MacConkey agar plates. One set of the culture plates was incubated aerobically, and the duplicate under microaerophilic conditions (5% CO₂). All plates were incubated at 37°C for 48 h. The isolates were subcultured for purity and identified using colony morphology, detection of hemolysis, Gram stain, and conventional biochemical tests (7).

Lung samples for virus isolation were stored in cryovials at -70°C and submitted to Institute of Virology, National Institute of Agriculture Technology, Morón, Buenos Aires, Argentina. Tissue samples were homogenized in Eagle's minimal essential medium (D-MEM) supplemented with 10% fetal bovine serum, and inoculated on MDBK (Madin-Darby Bovine Kidney) cells. Cell cultures were incubated at 37°C and 5% CO₂ for 5 days, and examined daily. After nine consecutive passages were carried out, cultures were tested for BRSV and BVDV by indirect fluorescent antibody (IFA) and PCR.

For histology and IHC, six lung samples obtained from the anterior, posterior cranial and caudal lobes of both left and right lungs were fixed in 10% buffered formalin, embedded in paraffin, sectioned at 3 µm, and

stained with both hematoxylin and eosin, and Gram stains. Selected paraffin-embedded tissues were submitted for immunohistochemical detection of *M. bovis* (Prairie Diagnostic Services Inc. Saskatoon Saskatchewan, Canada). Immunohistochemistry was conducted using a commercial staining platform (Ventana Medical System, Tucson AZ). Epitope retrieval was done by applying protease solution, and a mouse anti-*M. bovis* monoclonal antibody (mAb-5A10.2) was applied for 32 min at a dilution of 1: 800 (16).

Duplicate sections of each paraffin block were tested for *M. bovis* using a monoclonal antibody at 1:400 and 1:800 dilutions using positive and negative controls. No other *Mycoplasma* spp. were tested by IHC in this study.

Lung tissue of calves 1, 4 and, 5 were fixed in glutaraldehyde and post-fixed in osmium tetroxide for transmission electron microscopy (TEM). After fixation, tissues were dehydrated and embedded in epoxy resin. Semi-thin sections cut at 1-2 µm were mounted on slides and stained with toluidine blue. Slides were examined by light microscopy to select the areas of necrosis which were subsequently cut at 60 nm and contrasted with uranyl acetate and lead citrate. Thin sections were evaluated ultrastructurally with a JEM 1200 EX II (JEOL) transmission electron microscope.

Results

One of the affected calves in this study showed a clinical course that reached up to 59 days of illness. Table 1 summarizes farm locations, rates of infection, breed, age, and clinical signs of affected animals. There was no cytopathic effect observed in cell cultures, and all samples of lung tissue tested negative for BVDV and BRSV by IFA and PCR. *Trueperella pyogenes* was isolated from the lungs of two calves, *H. somni* from one calf, and *P. multocida* from another calf (Table 2).

On postmortem examination, the lungs showed cranioventral consolidation involving 40-70% of pulmonary parenchyma and the affected lung had a nodular texture and numerous focal to confluent white foci (Fig. 2). On the cut surface, the consolidated lung exhibited focal to confluent areas of necrosis composed largely of white, thick caseonecrotic exudate (Fig. 3).

Microscopically, the consolidated lung showed well-demarcated areas of necrosis surrounded by a thick layer of fibroblasts infiltrated with macrophages, lymphocytes, and few plasma cells (Figs. 4 and 5). These necrotic foci contained in the center abundant hyper eosinophilic and granular material (Fig. 6), while the periphery comprised bronchial walls some of which were dilated or effaced by the inflammatory reaction (Fig. 7). There was also thickening of the alveolar septa and suppurative bronchitis and bronchiolitis (Fig. 7), as well as hyperplasia of the bronchus-associated lymphoid tissue (BALT). The bronchiolar lumens of three calves were

Table 1. History of BRD outbreak, breed, age, clinical signs, and treatment information for animals with *Mycoplasma bovis* bronchopneumonia.

Farm, departament, province.	Calves no.	Month of outbreak / CI-M by BRD*	Breed/~age-months	Clinical signs/treatments† in animals autopsied.	FDO	DD
Dairy farm (A), Marcos Juárez, Córdoba.	1	May/4.5% (3/66) 1.5% (1/66)	Holstein /~3 m	Treated for BRD (1 tilm) but this calf did not respond to therapy and received additional treatments (2 tilm).	-	-
Feed lot (B), San Lorenzo, Santa Fe.	2	June/6.6% (12/182) 1.6 % (3/182)	Hereford/~8 m	Treated for BRD and severe claudication (1 tilm) but this steer did not respond and became markedly emaciated. Received additional treatment (1 tilm - 1 ceft).	18	77
	3	June/4% (4/100) 4% (4/100)	Aberdeen Angus/~6 m	Treated for BRD (1 tilm) and apparently improved but then was found dead in pen.	29	51
	4	April/12.5% (20/160) 5.6 % (10/160)	Crossbreed/~8 m	Treated for BRD (1 tilm) and apparently improved but then was found dead in pen.	15	26
Feed lot (C), Caseros, Santa Fe.	5	April/ 16% (40/250) 6% (15/250)	Aberdeen Angus/~7 m	First treated for BRD (1 tilm) and improved but then relapsed with BRD. Received additional treatment (1 tilm).	7	40
Feed lot (D), Calamuchita, Córdoba.	6	March/ 3.9 % (6/152) 1.3 % (2/152)	Crossbreed/~8 m	First treated for BRD (1 tilm) but this steer did not respond to therapy and received additional treatment (2 tilm).	13	50

ND, not done

BRD, bovine respiratory disease was determined by clinical examinations based on one or more of the following: depression (segregation from the group); persistent cough; excessive nasal discharge, dyspnea or hyperpnea, fever (above 40°C/104°F).

*CI, cumulative incidence; M, mortality.

†Tilmicosin (tilm): 10 mg/kg subcutaneously (SC) 3 days, Ceftiofur (ceft): 6.6 mg/KgSC 3 days.

FDO, fatal disease onset, DD, days of death.

obliterated by aggregates of fibroblast, collagen and newly formed blood capillaries (bronchiolitis obliterans) (Table 2). The lungs of calves 2, 3, 5, and 6 exhibited moderate pleural fibrosis. Calf 1 also had irregular and distinctive areas of coagulative necrosis surrounded by a rim of degenerated leukocytes with nuclear streaming resembling "oat-shaped cells" (Table 2). Viral inclusions were not observed in any of these lungs.

Immunohistochemistry showed abundant *M. bovis* antigen in the lungs of all six calves. The positive staining was observed mainly at the margin of the necrotic lesions, and to a lesser extent in the center of the necrotic foci (Fig. 8). *M. bovis* antigen was also detected in alveolar macrophages (Fig. 8).

Observation of lungs sections by electron microscopy revealed round to oval to pleomorphic structures with an approximated diameter ranging from 0.40 to 0.90 µm covered by a trilaminar membrane and varying electron density in the cytoplasm (Fig. 9). These structures were arranged in small aggregates or larger clusters throughout the caseonecrotic foci or admixed with the exudate present in the bronchial lumens (Fig. 9). These

structures were also observed primarily in the interstitial space between inflammatory cells such as lymphocytes, macrophages and neutrophils surrounding the necrotic areas.

Discussion

Pathological, ultrastructural, and immunostaining findings confirmed that the etiological agent incriminated in these six cases of chronic BRD in Argentina was *M. bovis*. To the best of our knowledge, this is the first report that unequivocally identifies *M. bovis* as the etiology of BRD in this country. Some researchers in Argentina had previously incriminated this organism as the putative agent in mastitis, but not related to pneumonia or respiratory disease (11).

The gross, microscopic and immunohistochemical findings in the lungs were identical to those previously reported in naturally occurring *M. bovis* pneumonia in other countries (5, 15, 20, 25, 26, 32). The same lesions have also been reproduced experimentally in calves (1, 2, 27).

Table 2: Gross findings, results of bacteriology and histological lung lesions in calves with natural infections with *Mycoplasma bovis*.

C no.	Necropsy findings	Bacteriology	Type 1 ²	Type 2 ³	Suppurative bronchitis and bronchiolitis	Obliterative bronchiolitis	BALT hyperplasia	Bronchointerstitial pneumonia
1	Severe subacute necro-suppurative bronchopneumonia, CNF and LNF 2-30 mm Ø, bronchiectasis.	<i>Trueperella pyogenes</i> .	+	++	+++	+++	++	++
2	Severe chronic necrotic bronchopneumonia, CNF 2-10 mm Ø, fibrous adhesions in pleura, arthritis in carpal joints, laminitis.	-	+++	-	+++	-	+++	+++
3	Subacute bronchopneumonia necro-suppurative, CNF and LNF 5-15 mm, fibrinous pleuritis.	<i>Trueperella pyogenes</i>	++	-	+++	+++	+	+++
4	Subacute bronchopneumonia fibrino-necrotic, CNF 2-5 mm Ø, interlobular edema.	<i>Histophilus somni</i>	+	-	++	-	++	+
5	Chronic severe necrotic bronchopneumonia, CNF 2-30 mm Ø, CNF involves the entire lobe, arthritis in carpal joints.	-	+++	-	+++	-	+++	+++
6	Chronic severe necrotic bronchopneumonia, CFN 2-5 mm diameter, fibrous adhesions in pleura.	<i>Pasteurella multocida</i>	+++	-	+++	+	+++	+++

CNF, foci of white-yellow, friable, caseous material surrounded by pale firm connective tissue.

LNF, foci of white-yellow, semi liquid material (purulent exudate).

-, none

²Type 1, caseonecrotic foci

³Type 2, coagulative necrotic foci

- None. + Mild. ++ Moderate. +++ Severe




Figure 1. Calf; moderate emaciation. This animal was exhibiting severe respiratory distress and had exudate in the nostrils. This calf also appeared emaciated.

Figure 2. Calf 5. Lung; bronchopneumonia; gross appearance. Lung shows cranioventral consolidation with a nodular texture. Inset: close-up of the pleural surface showing multiple white nodules.

Figure 3. Calf 5. Lung; cut surface; bronchopneumonia; gross appearance. Note numerous multifocal to coalescing raised nodules containing white caseous exudate. Bar =1.5 cm.

Figure 4. Calf 5. Lung, bronchopneumonia. Note numerous, well-demarcated foci of necrosis characterized by a large central core of necrotic hyper eosinophilic debris (asterisks). Hematoxylin and eosin, 2X.

Figure 5. Calf 5. Lung, bronchopneumonia. Note numerous, well-demarcated foci of necrosis characterized by a large central core of necrotic hyper eosinophilic debris (asterisk) surrounded by fibrous connective tissue. Hematoxylin and eosin, 5X.

Figure 6. Calf 5. Lung, bronchopneumonia; Close-up view of one necrotic focus showing abundant hyper eosinophilic granules (asterisk) surrounded by fibrous connective tissue infiltrated by mononuclear cells. Note some degenerated neutrophils at the margins of the hyper eosinophilic material (arrows). Hematoxylin and eosin, 40X. Inset: Aggregates of lymphocytes, plasma cells and macrophages.

Figure 7. Calf 5. Bronchiole filled with exudate composed of degenerated neutrophils and hyper eosinophilic granules (asterisk). The peribronchiolar tissues infiltrated with lymphocytes and plasma cells (arrows). Hematoxylin and eosin, 40X.

Figure 8. Calf. Lung, bronchopneumonia; immunohistochemistry. Note abundant positive antigen for *M. bovis* at the borders of the necrotic lesions (arrows). Inset: Positive staining in alveolar macrophages. Avidin-biotin peroxidase with hematoxylin counterstaining. 2X.

Figure 9. Transmission electron micrograph of many mycoplasma-like organisms aggregates between macrophages surrounding the caseonecrotic lesion. *Mycoplasma*-like organisms appear as pleomorphic oval structures with an intracellular material of varying electron density (asterisks) limiting membranes (white-arrowheads). Transmission electron microscopy 30,000X.

Feedlot systems appeared in Argentina around the 1990s and pneumonia has been the most common disease, as reported in the feedlot industry of North America (8, 22). The protracted clinical signs for all six cases in these BRD outbreaks in Argentina mimicked those for *M. bovis* pneumonia reported by others (26). *M. bovis* should always be suspected when a calf or steer has a history of an unresponsive BRD or when post mortem examination reveals bronchopneumonia with caseonecrotic nodules in the cranioventral lung lobes (9).

Although the caseonecrotic nodules resemble abscesses, these pulmonary structures are not pulmonary abscesses (*sensu stricto*), but rather segmentally distended bronchi filled with inspissated exudate (bronchiectasis) (8, 22). The pathogenesis of *M. bovis*-associated bronchiectasis is incompletely understood but presupposes long-term pulmonary inflammation that leads to the destruction of the bronchial walls (18, 22). This bronchial and bronchiolar changes are irreversible, and thus, explains why animals rarely respond to treatment and die, or are euthanized in extremis (5, 17, 26). Through experience and close observation, veterinarians and pathologists should be able to distinguish pulmonary abscesses frequently seen in cattle as an embolic sequel to liver abscesses, from caseonecrotic nodules and bronchiectasis, the long-term sequela of *M. bovis* bronchopneumonia. Nonetheless, it should also be noted that in some animals caseonecrotic nodules can progress into abscesses if there is a concurrent infection with *T. pyogenes* (10).

Bronchiolitis obliterans was another microscopic lesion seen in the *M. bovis* positive lungs of the calves in Argentina. This form of chronic obliterative bronchiolitis is a non-specific lesion resulting from severe injury and necrosis to the bronchiolar epithelium, where organized fibrovascular tissue progressively replaces the fibrin and neutrophils attached to the walls of the bronchioles (10, 22). The pathogenesis involved in bronchiolar necrosis and fibrosis in the bronchiolar lumen is still to be determined (18). Bronchiolitis obliterans is yet another contributing factor of respiratory distress in calves infected with *M. bovis* which is unlikely to resolve with antimicrobial treatment (17).

Calves 1 and 3 had Gram-positive coccobacilli within the *M. bovis*-IHC-positive caseonecrotic foci. This microscopic finding was not unexpected since co-infections with *T. pyogenes*, *P. multocida* and other mycoplasmas are common in natural and experimental *M. bovis* pneumonia (4, 6, 14, 15, 18).

The caseonecrotic nodules were also microscopically similar to those seen in *M. bovis* pneumonia elsewhere (10). Interestingly, the lungs of one calf (C1) showed not only hypereosinophilic granular lesions but also irregular areas of coagulative necrosis, a change most typically found with *M. haemolytica* infection (8, 22). There are two possible explanations for these co-existing microscopic findings: 1. *M. bovis* infection could have been a sequel or comorbidity to manheimiosis as

reported by others (22). 2. *M. bovis* could have caused in the same lung the two patterns of pulmonary necrosis as previously proposed by others (20).

All calves of the present study were unresponsive to antimicrobial treatment or suffered relapses of BRD which likely resulted from the combination of *M. bovis* capacity of evading the pulmonary defense mechanisms and immune responses (14, 15, 18) and that some strains of *M. bovis* are resistant to the antimicrobials used in feedlots to treat pneumonia, particularly tetracyclines and tilmicosin (3, 13, 28, 31). The same studies also suggested that tilmicosin was an ineffective therapeutic option for this agent, but extrapolating in vitro findings to the live animal conditions may not always be applicable.

Mycoplasma bovis is frequently found in the nasopharyngeal flora (10) and viral infections such as BoHV-1, BRSV, and PI-3 may predispose cattle to *M. haemolytica* infection, but the role of these viruses in the pathogenesis of *M. bovis* still needs elucidation. Since clinical signs of pneumonia appear at the late stages of *M. bovis* infection, it is difficult to know if there was a preceding viral infection in the calves of our study. This time difference between viral and *M. bovis* infections could also be the reason why the lungs of all calves in Argentina tested negative for respiratory viruses.

TEM and immunogold electron microscopy conducted in the lungs of experimentally infected cattle showed that *M. bovis* resides at the periphery of the necrotic foci, but it can also be found in the exudate within the bronchial lumen (21). The lungs from the affected animals in our study revealed comparable ultrastructural findings.

Polyarthritis is another condition frequently associated with *M. bovis* infection in cattle, and not surprisingly Calves 2 and 5 had in addition to pneumonia severe joint inflammation (1, 15). This joint infection is presumably a sequel to pneumonia where *M. bovis* disseminates hematogenously from the lung to the synovial membrane (23).

This first report should alert producers, veterinarians, and diagnosticians of the possible economic importance of *M. bovis* pneumonia in Argentina. Early detection and appropriate treatment are critical since this disease both clinically or silently progresses to irreversible lung damage and treating animals at the late stages of the diseases becomes expensive and futile. Further studies should determine the prevalence, morbidity, mortality and economic relevance of *M. bovis* pneumonia in the BRD complex in Argentina.

Acknowledgements

The authors would like to thank Gabriela Toledo, staff at the animals help group of the Agricultural Experiment Station Marcos Juárez, and Roxana Peralta of Central Electron Microscopy, School of Veterinary Sciences Service of National University of La Plata. The

work of Ms. Kathy Jones, Atlantic Veterinary College, is also appreciated.

This work was funded by project INTA PNSA-1115054, INTA AUDEAS CONADEV-940148 and ArgenINTA Foundation.

References

- Adegboye DS, Halbur PG, Nutsch RG, Kadlec RC, Rosembusch RF. *Mycoplasma bovis*-associated pneumonia and arthritis complicated with pyogranulomatous tenosynovitis in calves. *J Am Vet Med Assoc.* 1996;209:647-9.
- Arcangioli MA, Duet A, Meyer G, Dernburg A, Be'zille P, Poumarat F, LE GRAND D. The role of *Mycoplasma bovis* in bovine respiratory disease outbreaks in veal calf feedlots. *Vet J.* 2008;177:89-93.
- Ayling RD, Baker, SE, Peek ML, Simon AJ., Nicholas RAJ. Comparison of *in vitro* activity of danofloxacin, florfenicol, oxytetracycline, spectinomycin and tilmicosin against recent field isolates of *Mycoplasma bovis*. *Vet Rec.* 2000;146:745-7.
- Bell C.J, Blackburn P, Elliot M, Patterson TIAP, Ellison S, Lahuerta-marin A, Ball HJ. Investigation of polymerase chain reaction assays to improve detection of bacterial involvement in bovine respiratory disease. *J Vet Diagn Invest.* 2014;26:631-4.
- Booker CW, Abutarbush SM, Morley PS, Jim G.K, Pittman TJ, Schunicht OC, Perrett T, Wildman BK, Fenton RK, Guichon PT, Janzen ED. Microbiological and histopathological findings in cases of fatal bovine respiratory disease of feedlot cattle in Western Canada. *Can Vet J.* 2008;49:473-81.
- Bürki S, Frey J, Pilo P. Virulence, persistence and dissemination of *Mycoplasma bovis*. *Vet Microbiol.* 2015;179:15-22.
- Carter GR. Diagnostic procedures in veterinary bacteriology and mycology. 4th ed. Springfield: C.C. Thomas; 1984. 515 p.
- Caswell JL, Williams KJ. Respiratory system. In: Maxie MG, editor. *Jubb, Kennedy and Palmer's Pathology of Domestic Animals.* 6th ed., vol. 2. St. Louis: Elsevier; 2016. p. 465-591.
- Caswell JL, Archambault M. *Mycoplasma bovis* pneumonia in cattle. *Anim Hlth Res Rev.* 2008;8:161-86.
- Caswell JL, Bateman KG, Hugh YC, Castillo-Alcala F. *Mycoplasma bovis* in respiratory disease of feedlot cattle. *Vet Clin North Am Food Animal Pract.* 2012;26:365-79.
- Cerdá R, Xavier J, Sansalone P, De La Sota R, Rosembusch R. Aislamiento de *Mycoplasma bovis* a partir de un brote de mastitis bovina en una vaquería de la provincia de Buenos Aires. Primera comunicación en la República Argentina. *Rev La Am Microbiol.* 2001;42:7-11.
- Ellis JA. The immunology of the bovine respiratory disease complex. *Vet Clin North America Food Anim Pract.* 2001;17:535-50.
- Francoz D, Fortin M, Fecteau G, Messier SE. Determination of *Mycoplasma bovis* susceptibilities against six antimicrobial agents using the E test method. *Vet Microbiol.* 2005;105:57-64.
- Fulton, RW, Blood SK, Panciera RJ, Payton ME, Ridpath JF, Confer AW, Saliki JT, Burge TL, Welsh RD, Johnson BJ, Reck A. Lung pathology and infectious agents in fatal feedlot pneumonias and relationship with mortality, disease onset, and treatments. *J Vet Diagn Invest.* 2009;21:464-77.
- Gagea MI, Bateman KG, Van Dreumel T, Macewen BJ, Carman S, Archambault M, Shanahan RA, Caswell JL. Diseases and pathogens associated with mortality in Ontario beef feedlots. *J Vet Diagn Invest.* 2006;18:18-28.
- Haines DM, Chelack BJ. Technical considerations for developing enzyme immunohistochemical staining procedures on formalin-fixed, paraffin-embedded tissues for diagnostic pathology. *J Vet Diagn Invest.* 1991;3:101-12.
- Haines DM, Martin KM, Clark EG, Jim J, Janzen ED. The immunohistochemical detection of *Mycoplasma bovis* and bovine viral diarrhoea virus in tissues of feedlot cattle with chronic, unresponsive respiratory disease and/or arthritis. *Can Vet J.* 2001;42:857-60.
- Hermeyer K, Buchenau I, Thomasmeyer A, Baum B, Spersger J, Rosengarten R, Hewicker-Trautwein M. Chronic pneumonia in calves after experimental infection with *Mycoplasma bovis* strain 1067: characterization of lung pathology, persistence of variable surface protein antigens and local immune response. *Acta Vet Scand.* 2012;54:1-11.
- Horwood PF, Schibrowski ML, Fowler EV, Gibson JS, Barnes TS, Mahony TJ. Is *Mycoplasma bovis* a missing component of the bovine respiratory disease complex in Australia? *Aust Vet J.* 2014;92:185-91.
- Khodakaram-Tafti A, López A. Immunohistopathological findings in the lungs of calves naturally infected with *Mycoplasma bovis*. *J Vet Med A Physiol Pathol Clin Med.* 2004;51:10-4.
- Kleinschmidt S, Spersger J, Rosengarten R, Hewicker-Trautwein M. Long-term survival of *Mycoplasma bovis* in necrotic lesions and in phagocytic cells as demonstrated by transmission and immunogold electron microscopy in lung tissue from experimentally infected calves. *Vet Microbiol.* 2013;162:949-53.
- López A, Martinson SA. Respiratory System, Mediastinum, and Pleurae. In: Zachary JF, editor. *Pathologic Basis of Veterinary Diseases.* 6th ed. St. Louis: Elsevier; 2017. p. 471-560.
- Nicholas RAJ, Ayling RD. *Mycoplasma bovis*: disease, diagnosis, and control. *Res Vet Sci.* 2003;74:105-12.

24. Panciera R, Confer AW. Pathogenesis and pathology of bovine pneumonia. *Vet Clin North America Food Anim Pract.* 2010;26:191-214.
25. Radaelli E, Luini M, Loria GR, Nicholas RAJ, Scanziani E. Bacteriological, serological, pathological and immunohistochemical studies of *Mycoplasma bovis* respiratory infection in veal calves and adult cattle at slaughter. *Res Vet Sci.* 2008;85:282-90.
26. Ramírez-Romero R, Chavarría-Martínez B, Nevárez-Garza, AM, Rodríguez-Tovar LE, Dávila-Martínez C, Hernández-Vidal G, Hernández-Escañero JJ, López-Mayagoitia A. Demostración histoquímica de *Mycoplasma bovis* en lesiones neumónicas crónicas en ganado de corral de engorda. *Vet Mex.* 2010;41:289-96.
27. Rodríguez F, Bryson DG, Ball HJ, Forster F. Pathological and immunohistochemical studies of natural and experimental *Mycoplasma bovis* pneumonia in calves. *J Comp Pathol.* 1996;115:151-62.
28. Rodríguez F, Kennedy S, Bryson TD, Fernandez A, Rodríguez JL, Ball HJ. Immunohistochemical method of detecting *Mycoplasma* species antigen by use of monoclonal antibodies on paraffin sections of bovine and caprine lungs. *Zentralbl Veterinarmed B.* 1996;43:429-38.
29. Rosenbusch, RF, Kinyon JM, Apley M, Funk ND, Smith S, Hoffman JL. *In vitro* antimicrobial inhibition profiles of *Mycoplasma bovis* isolates recovered from various regions of the United States from 2002 to 2003. *J Vet Diagn Invest.* 2005;17:436-41.
30. Terlaak EA, Wentink GH, Zimmer GM. Increased prevalence of *Mycoplasma bovis* in the Netherlands. *Vet Quart.* 1992;14:100-4.
31. Thomas A, Nicolas C, Dizier I, Mainil J, Linden, A. Antibiotic susceptibilities of recent isolates of *Mycoplasma bovis* in Belgium. *Vet Rec.* 2003;153:428-31.
32. Yilmaz R, Cangul IT, Onat K, Akkoc A, Ozyigit MO, Akdesir E. Histopathological, immunohistochemical and bacteriological characterization of *Mycoplasma bovis* pneumonia in cattle. *Pakistan Vet J.* 2016;36:316-21.