Braz J Vet Pathol, 2025, vol. 18, Suppl. 1, e018S107 DOI: https://doi.org/10.24070/bjvp.1983-0246.018S107





Diagnostic Exercise From the Latin Comparative Pathology Group and the Davis-Thompson Foundation

Small intestinal entrapment through a mesovarium ligament rent in a mare

Ji Ann Greenberg¹, Sydney Hoff¹, Raquel R Rech¹*

Department of Veterinary Pathobiology, College of Veterinary Medicine and Biomedical Sciences, Texas A&M University, College Station, TX, USA.

*Corresponding author: rrech@cvm.tamu.edu

History:

An 18-year-old, American Paint mare presented for acute colic for a suspected colon torsion. On presentation, the blood and abdominal lactate were 1.3 mmol/L (normal range 0-2 mmol/L) and 5.3 mmol/L (normal range <2 mmol/L), respectively. In the following morning, her blood and abdominal lactate increased to 7.3 mmol/L and 9.9 mmol/L. Due to uncontrollable pain, lack of a surgical option due to financial limitations, and increasing suspicion for a strangulating lesion, euthanasia was elected.

Necropsy findings:

An approximately 2.4 m segment of ileum and distal jejunum was diffusely dark red and dilated, starting at the ileocecal junction, and extending orally. The orad jejunal loops were diffusely dilated up to 8 cm (Figure 1). The dark red intestinal loops were entrapped by a thin band of tissue formed by a 6 cm in diameter in the mesovarium ligament (Figures 2 and 3). The entrapped segment contained hemorrhagic, watery contents, and the mucosa was roughened and covered by fibrin. The mesenteric vessels were also congested and had marked venous thrombosis.

Follow-up questions:

- Morphologic diagnosis
- *Differential diagnoses* (Figure 1)



*The Diagnostic Exercises are an initiative of the Latin Comparative Pathology Group (LCPG), the Latin American subdivision of The Davis-Thompson Foundation (DTF). These exercises are contributed by members and non-members from any country of residence.

Consider submitting an exercise! A final document containing this material with answers and a brief discussion will be posted on the DTF website:

https://davisthompsonfoundation.org/diagnostic-exercise/

Editor-in-chief for this Diagnostic Exercise: Claudio Barros Associate Editor for this Diagnostic Exercise: Raquel Rech



Braz J Vet Pathol, 2025, vol. 18, Suppl. 1, e018S107 DOI: https://doi.org/10.24070/bjvp.1983-0246.018S107




Figure 1. Entrapped congested jejunal segments with venous thrombosis.



Figure 2. Close up view of the segment of distal jejunum entrapped by a thin band of tissue (rent).

ANSWERS

Morphologic diagnosis:

Ileal-jejunal incarceration through a right mesovarium ligament rent with acute venous congestion and thrombosis.

Differential diagnoses (Figure 1):

Other possible causes for segmental small intestinal strangulation in horses include pedunculated mesenteric lipoma, epiploic foramen entrapment, mesenteric rent, gastrosplenic ligament entrapment, volvulus/torsion, and inguinal/scrotal/umbilical/diaphragmatic hernias, mesodiverticular band entrapment, rents in proximal aspect of the cecocolic fold, and mesometrial tears.

Comments:

Colic is one of the most common emergencies in equine practice, and a frequent cause is small intestinal

strangulation. Strangulating obstruction results from occlusion of the intestinal lumen and its blood supply causing not only intraluminal distention, but also ischemic injury. The degree of injury can be highly variable and is dependent on the nature of the vascular occlusion. Veins have thinner, more compliant walls, and lower hydrostatic pressure than arteries and consequently are typically occluded first. In the early stages, there is an absence of patent outflow due to the venous occlusion but continued delivery of arterial blood, resulting in a hemorrhagic strangulating obstruction, characterized by ischemic injury and tissue congestion. If the strangulating obstruction exerts enough pressure and occludes both the veins and arteries, it is termed an ischemic strangulating obstruction, resulting in rapid degeneration of the mucosa (3). The ischemic injury eventually leads to devitalization of the tissue and is associated with a poor prognosis, especially without surgical treatment.

Common causes of small intestinal strangulation include pedunculated lipoma, epiploic foramen entrapment, mesenteric rents, volvulus/torsion, gastrosplenic ligament entrapment, and inguinal/scrotal/umbilical hernias. Rare causes of intestinal strangulation are mesodiverticular bands (1), rents in proximal aspect of the cecocolic fold (7), which is the result of a persistent vitelline artery (1), and mesometrial rents (8). The mesometrium is the mesentery of the uterus that forms the broad ligament of the uterus, which is a peritoneal fold that attaches the uterus to the abdominal and pelvic walls. The broad ligament is subdivided into the mesosalpinx, mesovarium and mesometrium (4). To date, there have been two reports of small intestinal strangulation through a rent in the mesometrium in pregnant mares (2,6), one in a nonpregnant mare (9) and one in a mare that did not disclose pregnancy status (10). In women, defects in the broad ligament leading to intestinal incarceration are exceedingly rare. It usually presents as right-sided rents, with the most common herniating organ being the ileum, although cases involving the colon or ovary have been reported. There is an increase in reports in multiparous women, patients who have undergone the Webster-Baldy operation, and patients with a history of salpingitis or endometriosis. A congenital etiology should be considered in women that are nulliparous or who never underwent an abdominal surgical procedure (5).

The etiology of the mesovarium rent in this case is unknown, but highest priority is given to previous trauma (acquired) or a congenital defect. Possible traumatic causes of an acquired defect in the ligament include past surgery (iatrogenic), previous foaling especially if there were complications (dystocia), any manipulation of the uterus, a direct tear secondary to trauma to the area, or weakened tissues. Unfortunately, a complete medical history of this mare was not provided. This case aims to demonstrate that although rare, a mesovarium ligament rent should be a differential for strangulating obstructions in mares.

Braz J Vet Pathol, 2025, vol. 18, Suppl. 1, e018S107 DOI: https://doi.org/10.24070/bjvp.1983-0246.018S107




Figure 3a-c. The thin band of tissue is close to the right ovary (arrow) (a) that creates a 6 cm in diameter defect (b) in the mesovarium ligament (c).

References:

- 1. Abutarbush SM, Shoemaker RW, Bailey JV. Strangulation of the small intestines by a mesodiverticular band in 3 adult horses. Can Vet J. 2003;44(12): 1005-1006.
- 2. Becht JL, McIlwraith CW. Jejunal displacement through the mesometrium in a pregnant mare. Journal of the American Vet Medical Association 1980; 177(5): 436.
- 3. Blikslager AT. Pathophysiology of Gastrointestinal Obstruction and Strangulation. In: The Equine Acute Abdomen. 3rd ed. Editors A.T. Blikslager, N.A. White, J.N. Moore and T.S. Mair. Whiley; 2017. p. 102-104.
- Craig ME, Sudanagunta S, Billow M. Anatomy, Abdomen and Pelvis: Broad Ligaments. In StatPearls. Stat-Pearls Publishing; 2023.
- 5. Demir H, Scoccia B. Internal Herniation of Adnexa Through a Defect of the Broad Ligament: Case Report and Literature Review. Journal of Minimally Invasive Gynecology 2010; 17(1): 110-112.

- Freeman DE, Schaeffer DJ. Clinical comparison between a continuous Lembert pattern wrapped in a carboxymethylcellulose and hyaluronate membrane with an interrupted Lembert pattern for one-layer jejunojejunostomy in horses. Equine Vet Journal 2011; 43(6): 708-713.
- 7. Gayle JM, Macharg MA, Smallwood JE. Strangulating Obstruction Caused by Intestinal Herniation Through the Proximal Aspect of the Cecocolic Fold in 9 Horses. Vet Surgery 2001; 30: 40-43.
- 8. Schaffer PA, Sonis JM, Rodriguez JS, Han S. Case report: Ileo-jejunal entrapment in a uterine mesometrium rent of a non-pregnant mare. Clinical Theriogenology 2012; 4(1): 55-6.
- 9. Steward SK, Bauck AG, Zoll W, Conway JA, Freeman DE. Small intestinal strangulation in a tear in the mesometrium of a nonpregnant mare. Equine Vet Education 2017; 30(11): 591-593.
- Tennant B, Wheat JD, Meagher DM. Observations on the causes and incidence of acute intestinal obstruction in the horse. Proc. Am. Assoc. Equine Practice 1975; 21:426-438.