



## Diagnostic Exercise

From the Latin Comparative Pathology Group and the Davis-Thompson Foundation

# Bursal cryptosporidiosis in a kelp gull (*Larus dominicanus*)

Diego A. Yañez<sup>1\*</sup>

Laboratorio AnimaLab Diagnóstico, Quilpué, Chile.


Corresponding author: [animalabdiagnostico@gmail.com](mailto:animalabdiagnostico@gmail.com)

### History:

A juvenile kelp gull (*Larus dominicanus*) was found dead.

### Necropsy findings:

The bird was in good body condition. Gross lesions were restricted to the head, with extensive skull fractures, severe intracranial and subcutaneous hemorrhage (*in vivo* trauma). Samples were collected for histopathology (Figs.1-4).



**Figures: 1-4.** Cloacal bursa. There is moderate bursal epithelial hyperplasia. Abundant, small, round basophilic protozoa morphologically consistent with *Cryptosporidium* spp. are lining the apical portion of the epithelium. Minimal intraepithelial heterophils are seen. Lumina are distended by a small amount of sloughed epithelial cells, heterophils, and cell debris. No changes in lymphoid tissue are noted.



*\*The Diagnostic Exercises are an initiative of the Latin Comparative Pathology Group (LCPG), the Latin American subdivision of The Davis-Thompson Foundation (DTF). These exercises are contributed by members and non-members from any country of residence. Consider submitting an exercise! A final document containing this material with answers and a brief discussion will be posted on the DTF website: <https://davis-thompsonfoundation.org/diagnostic-exercise/>*



*Editor-in-chief for this Diagnostic Exercise: Claudio Barros  
Associate Editor for this Diagnostic Exercise: Daniel Rissi*

**Follow-up questions:**

- *Morphologic diagnosis*
- *Etiology*
- *Name other organs that can be affected by this agent, in avian species*

**ANSWERS****Histologic description:**

Cloacal bursa (Figs. 1-4): There is moderate bursal epithelial hyperplasia. Abundant, small, round basophilic protozoa morphologically consistent with *Cryptosporidium* spp. are lining the apical portion of the epithelium. Minimal intraepithelial heterophils are seen. Lumina are distended by a small amount of sloughed epithelial cells, heterophils, and cell debris. No lymphoid tissue changes are noted.

**Morphologic diagnosis:**

Cloacal bursa: Minimal, acute heterophilic bursitis with epithelial hyperplasia and surface-associated protozoa.

**Etiology:**

*Cryptosporidium* spp. (most commonly *C. baileyi*).

**Comments:**

*Cryptosporidium* is an apicomplexan protozoan that infects birds, mammals, fish, reptiles, amphibians, and humans, with 19 known zoonotic and potentially zoonotic species (4,5,8). *Cryptosporidiosis* can be defined as a secondary and no host-specific parasitism, more often producing clinical disease in immunocompromised individuals (4,5,8).

The most described species of *Cryptosporidium* infecting birds are *C. baileyi*, also being the most reported in laridae (3), *C. meleagridis*, mostly reported in turkeys, and *C. galli* in chickens (4,5), although different parasite species can infect different hosts, including anatidae, accipitridae, falconidae, passeridae, phasianidae, psittacidae and tytonidae (8).

*Cryptosporidium* can infect nasal the cavity, trachea, bronchi, and lungs, producing epiphora, conjunctival redness, swollen eyes, cough, and abundant mucous secretion accumulation in the airways (4,5). Gastrointestinal infections are reported in the proventriculus, intestine, colon, cloaca (1,5,6), and bursa (1,5,7) of chickens and turkeys, leading to diarrhea and weight loss (6). The urinary tract can also be infected, producing weight loss and

systemic gout (2). The most common pathologic features of *Cryptosporidium* infection are the presence of 2 to 8 µm round protozoan parasites located in the apical portion of the epithelium, epithelial hyperplasia and sloughing of cell debris are mixed with mucous exudate. Mixed heterophilic and lymphoplasmacytic infiltrates can be detected in the lamina propria of the affected organs (1). *C. baileyi* is the most reported agent in laridae, producing respiratory cryptosporidiosis in *Chroicocephalus rubicundus* and *Larus argentatus* young gulls, but subclinical infections of the cloacal bursa and cloaca are most commonly reported (3).

The definitive diagnosis of cryptosporidiosis can be achieved by immunohistochemistry, polymerase chain reaction (PCR), and electron microscopy showing the characteristic intracellular and epicytoplasmic distribution of the protozoa (5).

There is limited information about cryptosporidiosis in *Larus dominicanus* and its role in the dissemination of this agent in the environment is unknown.

**References**

1. Abdul-Aziz T. Avian histopathology, 4th ed. Jacksonville: American Association of Avian Pathologists; 2016. p 45-46.
2. Curtiss, J. B., Leone, A. M., Wellehan, J. F. X., Emerson, J. A., Howarth, E. W., Farina, L. L. Renal and cloacal cryptosporidiosis (*Cryptosporidium avian* genotype V) in a Major Mitchell's cockatoo (*Lophochroa leadbeateri*). Journal of Zoo and Wildlife Medicine, 2015 46(4), 934-937.
3. Hubálek, Z. Pathogenic microorganisms associated with gulls and terns (*Laridae*). Journal of Vertebrate Biology 2021 70.3:21009-1.
4. Lindsay DS, Blagburn BL. Cryptosporidiosis in birds, 133-148. In: Dubey, J.P., Speer, CA., Fayer R., Eds., Cryptosporidiosis in Man and Animals, CRC Press, Boca Raton; 1990.
5. Nakamura, A and Vasconcelos Meireles, M. Cryptosporidium infections in birds-a review. Revista Brasileira de Parasitologia Veterinária 2015 24.3: 253-267.
6. Ravich ML, Reavill, DR, Hess, L, Childress AL, Wellehan Jr, JF Gastrointestinal cryptosporidiosis in captive psittacine birds in the United States: a case review. Journal of Avian Medicine and Surgery, 2014 297-303.
7. Schulze, C., Kämmerling, J., Kutzer, P., & Richter, B. *Cryptosporidium baileyi* Infection in red-breasted merganser ducklings. Journal of Comparative Pathology, 2012 146(1), 70.
8. Wang Y., Zhang K, Chen Y, Li X, Zhang L. Cryptosporidium and cryptosporidiosis in wild birds: A One Health perspective. Parasitology Research, 2021 120(9), 3035-3044.