

1 **Case Report**

2
3 **Fatal canine metaldehyde poisoning in Southern Brazil**

4
5 Rômulo Debortoli¹  (<https://orcid.org/0000-0002-0077-5987>) Alexandre Arenales^{1*} 
6 (<https://orcid.org/0009-0006-5411-6983>)

7
8 ¹ Universidade do Oeste de Santa Catarina (Unoesc), Unidade de Atendimento Veterinário. Campos
9 Novos, SC, Brazil.

10 ² Veterinarian, Alô Vet - Atendimento Veterinário, Campos Novos, SC, Brazil

11
12 ***Corresponding author:** alexandre.arenales88@gmail.com

13
14 Submitted: August 7th, 2025. Accepted: December 19th, 2025.

15
16 **Abstract**

17 Metaldehyde is commonly used as a molluscicide, and accidental ingestion in dogs is a
18 recognized toxicologic emergency with potentially fatal outcomes. A 4-month-old female Pit Bull
19 was presented to a veterinary emergency after ingesting an unknown amount of a 5% metaldehyde-
20 based slug bait (Metarex®). Clinical signs began within minutes and included stupor, hypersalivation,
21 and generalized tonic-clonic seizures. Supportive care was provided, including anesthetic coma with
22 propofol and isoflurane, gastric lavage, diazepam, N-acetylcysteine, and oxygen therapy; however,
23 the animal died in eight hours. At necropsy, the stomach and intestines contained abundant blue-to-
24 green amorphous and finely granular material, as well as plastic fragments consistent with Metarex®
25 packaging. No significant microscopic alterations were observed in internal organs. The diagnosis
26 was based on the clinical course, history of exposure, and characteristic gastrointestinal contents. In

27 suspected cases, identification of metaldehyde in the gastrointestinal tract can aid diagnosis when
28 toxicologic confirmation is unavailable.

29

30 **Keywords:** molluscicide, slug bait, acute poisoning.

31

32 **Introduction**

33

34 Metaldehyde is a cyclic tetramer of acetaldehyde, widely used as a molluscicide to control
35 terrestrial gastropods (slugs and snails), such as *Achatina fulica*. Commercial baits are typically
36 formulated as blue-to-green granules or pellets with concentrations ranging from 1.5% to 8% and act
37 by inducing dehydration and the death of mollusks (1, 3, 7). Bran or molasses are sometimes added
38 to the pellets, which increases bait palatability, particularly for dogs (4). In dogs, the estimated oral
39 median lethal dose (LD₅₀) of metaldehyde ranges from 100 to 1000 mg/kg, although a commonly
40 cited value is around 210 mg/kg, based on older experimental studies (1, 3, 7).

41 The precise mechanism of toxicity in mammals remains unclear. In murine models, it is
42 considered a neurotoxic agent that inhibits neurotransmitters such as serotonin and norepinephrine,
43 and primarily gamma-aminobutyric acid (GABA), leading to increased monoamine oxidase activity,
44 a lowered seizure threshold, and neuronal excitation (5). Metaldehyde poisoning in dogs can be
45 rapidly fatal. Clinical signs usually appear within 30 minutes to 3 hours and include neurological and
46 systemic signs such as convulsion, hypersalivation, tremors, hyperesthesia, seizures, hyperthermia,
47 ataxia, vomiting, metabolic acidosis, tachypnea, dyspnea, hyperventilation, and respiratory
48 depression (1, 5).

49 There is currently no specific antidote for metaldehyde poisoning, thus the treatment is
50 supportive until the toxin is eliminated (3). Despite early intervention, prognosis remains uncertain
51 in severe cases; and reported survival rates in treated dogs vary: in one study, the survival rate was

52 81% (5), and in another, 83% (7). This case report describes the clinical progression, pathological
53 findings, and fatal outcome of acute metaldehyde poisoning in a dog from Southern Brazil.

54

55 **Case description**

56

57 A 4-month-old female Pit Bull was presented to a veterinary emergency service in Campos
58 Novos, Santa Catarina, Brazil, shortly after accidental ingestion of a commercial molluscicide
59 containing metaldehyde (Metarex®). The owner found that the dog had torn open the slug bait
60 package and ingested part of its contents; the product and packaging were documented and referred
61 (Fig. 1).

62 Blue to green amorphous content, residue of metaldehyde pellets, was visibly present in the
63 oral cavity, and the dog exhibited stupor, sialorrhea, and generalized tonic-clonic seizures (S1). Rectal
64 temperature was 37.8 °C, heart rate 165 bpm, respiratory rate 36 breaths per minute, and systolic
65 arterial pressure 140 mmHg. Anesthetic coma was induced using propofol and maintained with
66 isoflurane. Gastric lavage was performed via nasogastric intubation, and the treatment protocol
67 included diazepam (1 mg/kg IV), fluid therapy (60 mL/kg/h), N-acetylcysteine (150 mg/kg IV),
68 omeprazole (1 mg/kg), and oxygen. During the 8 hours of anesthetic coma, the animal experienced
69 progressive cardiovascular and respiratory depression, hypotension, bradycardia, and decreased
70 respiratory rate and body temperature, leading to death.

71 Macroscopically, the stomach lumen was filled with a large amount of blue to green amorphous
72 and finely granular content (Fig. 2), along with pieces of plastic compatible with the slug bait packet
73 referred by the owner (Fig. 3). Similar content was found in the lumen of all segments of the small
74 intestine (Fig. 4) and the initial portion of the colon. Additionally, there was a focal extensive area of
75 edema and hemorrhage in the subcutaneous tissue of the lateral thoracic region, attributed to
76 prolonged lateral decubitus. No significant microscopic alterations were found in internal organs.

77

78 **Discussion**

79

80 This report describes a case of fatal metaldehyde poisoning in a four-month-old Pit Bull, with
81 diagnosis supported by the presence of blue to green amorphous and finely granular gastrointestinal
82 contents and fragments of the commercial molluscicide packaging. The giant African snail (*Achatina*
83 *fulica*) is a well-known invasive species in Southern Brazil (6), and the use of metaldehyde-based
84 molluscicides is a common control strategy. Although not previously published in the region, cases
85 of metaldehyde poisoning in dogs are not unusual in the clinical routine of small animal practitioners,
86 who are generally aware of this risk.

87 Although the ingested amount was not precisely measured, the animal had access to a full 200
88 g package containing 5% metaldehyde (Metarex®) (Fig.1) (7), likely exceeding the lethal dose.
89 Moreover, the difficulty in determining the exact amount ingested has also been highlighted
90 previously (1). Clinical signs in this case were consistent with those previously described (1, 5),
91 including seizures, hypersalivation, and neurologic deterioration. The rapid onset of seizures and
92 death reflects the fulminant course sometimes observed in high-dose exposures (3, 5). No other gross
93 lesions were identified, apart from the blue-to-green content in the gastrointestinal tract, which,
94 together with the clinical history, was sufficient to establish the diagnosis. Nonspecific lesions such
95 as thoracic cavity hemorrhages have been described in other reports (2), but were not observed in this
96 case.

97 In conclusion, necropsy findings are a valuable diagnostic tool for metaldehyde poisoning in
98 dogs. Identification of bait-like material in gastrointestinal contents can support diagnosis when
99 laboratory confirmation is unavailable.

100

101 **Supplementary Material**

102 The online version contains supplementary material available at
103 <https://doi.org/10.24070/bjvp.1983-0246.019002>.

104

105 **Conflict of Interest**

106 The authors declare no competing interests.

107

108 **References**

109

- 110 1. Bates NS, Sutton NM, Campbell A. Suspected metaldehyde slug bait poisoning in dogs: a
111 retrospective analysis. *Vet Rec.* 2012;171(14):351. doi: 10.1136/vr.100734.
- 112 2. Botelho AFM, Machado AMD, da Silva RHS, Costa FS, Carvalho RM, Martins MI. Fatal
113 metaldehyde poisoning in a dog confirmed by gas chromatography. *BMC Vet Res.*
114 2020;16:139.
- 115 3. De Roma A, Miletti G, D'Alessio N, Rossini C, Vangone L, Galiero G, et al. Metaldehyde
116 poisoning of companion animals: a three-year retrospective study. *J Vet Res.* 2017;61(3):307–
117 11. doi: 10.1515/jvetres-2017-0041.
- 118 4. Dolder LK. Metaldehyde toxicosis. *Vet Med.* 2003;98:213-215.
- 119 5. Dutil GF, Berny P. A prospective study on clinical signs, management, outcomes, and delayed
120 neurologic sequelae due to metaldehyde poisoning in 26 dogs. *Open Vet J.* 2023;13(5):510-4.
121 doi: 10.5455/OVJ.2023.v13.i5.2.
- 122 6. Thiengo SC, Faraco FA, Salgado NC, Cowie RH, Fernandez MA. Rapid spread of an invasive
123 snail in South America: the giant African snail, *Achatina fulica*, in Brazil. *Biol Invasions.*
124 2007;9(5):693-702. doi:10.1007/s10530-006-9069-6.
- 125 7. Yas-Natan E, Segev G, Aroch I. Clinical, neurological and clinicopathological signs, treatment
126 and outcome of metaldehyde intoxication in 18 dogs. *J Small Anim Pract.* 2007;48(8):438-43.
127 doi: 10.1111/j.1748-5827.2007.00360.x.

128

129



130

131 **Figure 1.** A package of metaldehyde-based slug bait (Metarex®) referred by the owner, opened and
132 thorned by the dog. Characteristic blue pellets are visible on the left side.

133



134

135 **Figure 2.** Dog, metaldehyde poisoning, stomach with abundant blue to green amorphous and finely
136 granular content compatible with digested metaldehyde-based slug bait.

137



138

139 **Figure 3.** Dog, metaldehyde poisoning, stomach content compatible with the package of
140 metaldehyde-based slug bait (Metarex®) shown in Figure 1.

141



142

143 **Figure 4.** Dog, metaldehyde poisoning, small intestine with blue to green amorphous and finely
144 granular content compatible with digested metaldehyde-based slug bait.

145

146