



Original Full Paper

Caffeine exposure causes downregulation of the expression of genes related to osteogenesis and chondrogenesis in Zebrafish (*Danio rerio*)

Bruno Machado Bertassoli¹ , Gilcinea de Cassia Santana¹ , Julia Meireles Nogueira² ,
Rayan Silva de Paula² , Erika Cristina Jorge² , Helder de Oliveira Guilherme¹ ,
Endrigo Gabellini Leonel Alves³ , Amanda Maria Sena Reis² , Rogéria Serakides¹ , Natália Melo Ocarino^{1,*} 

¹Escola de Veterinária, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

²Instituto de Ciências Biológicas da Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

³Universidade de Uberaba (UNIUBE), Uberaba, Minas Gerais, Brazil

*Corresponding author: nataliaocarino@gmail.com

Submitted: September 25th, 2023. Accepted: November 21st, 2023.

Abstract

With the aim of evaluating the effects of caffeine on the expression of gene transcripts related to osteogenesis and chondrogenesis, during zebrafish embryonic development the embryos were divided into five groups: control (without caffeine), 0.25 mM caffeine, 0.5 mM caffeine, 1.0mM caffeine, and 2.0mM caffeine. Embryos were placed in cell culture plates with aquarium water at 28°C for 72 hours post-fertilization (hpf). Embryonic development was followed at times of six, 12 hpf, 24 hpf, 30 hpf, 48 hpf, 54 hpf, and 72 hpf. At 72 hpf, the expression of Sox9a, runx2b, osteocalcin, osteopontin, collagen 2a1, and bmp2a was analyzed using real-time RT-PCR. The 0.25mM caffeine group showed embryonic development that was visually similar to the control group despite having a significantly lower relative expression of bmp2a, runx2b, and collagen 2a1. In the 0.5mM and 1mM caffeine groups, morphological alterations such as tail folding, pericardial edema, and yolk sac deformation were observed, and the relative expression levels of bmp2a, runx2b, and collagen 2a1 were significantly lower. In contrast, osteocalcin expression was significantly higher than in the control group. In the 2mM caffeine group, the embryos did not hatch until 72 hpf and showed tail docking and pericardial and yolk sac edema. The relative expression of bmp2a and runx2b was significantly lower, whereas that of osteocalcin, osteopontin, and collagen 2a1 did not differ from that of the control group. We conclude that caffeine altered the expression of gene transcripts related to osteogenesis and chondrogenesis in zebrafish embryos, even in embryos without visible morphological changes.

Keywords: 1,3,7-trimethyl xanthine, contaminants, embryos, fish, morphology

Introduction

Pharmaceuticals, personal care products, and industrial chemicals are being increasingly identified in aquatic environments worldwide. They have aroused increasing concern about preserving aquatic ecosystems because they can damage ecosystems, even at residual levels (21). Caffeine

(1,3,7-trimethyl xanthine) is one of the most emerging contaminants in natural aquatic environments and is an indicator of anthropogenic contamination. It is an alkaloid present in coffee, tea, soft drinks, beverages, and various pharmaceutical and personal care products. Caffeine can contaminate freshwater and saltwater ecosystems through sewage, improper disposal of drugs, and medical waste. Although wastewater

treatment is efficient for caffeine removal, it is insufficient to prevent caffeine from being transported through aquatic systems because of its relative stability under different environmental conditions and significant and increasing consumption (34, 62).

In freshwater ecosystems, caffeine concentrations have been found in concentrations of up to 1500 ng/L. The highest concentration detected in marine environments was 11,000 µg/L in the Port of Darwin, Northern Australia (25). In US marine waters, caffeine concentrations range from 2 to 5860 ng/L (62). In Europe, the highest concentrations of caffeine were found in the Aegean Sea (3068 ng/L) (39) and the Arade River in Portugal (600–804 ng/L) (27). Caffeine has also been identified in marine environments in several other countries and various aquatic organisms, including corals, algae, mussels, and fish (62).

Research has shown that caffeine present in aquatic environments can influence the health of organisms, affecting the reproduction and survival of species and, consequently, compromising the food chain of ecosystems. Caffeine concentrations similar to those found in natural ecosystems cause adverse effects in several aquatic species maintained under laboratory conditions (62). Embryos of *Palaemonetes pugio* shrimp exposed to 20 mg/L caffeine for five days showed delayed development and egg hatching (26). The effect of caffeine exposure (0.5–18 µg/L) on the regenerative capacity of the polychaete *Diopatra neapolitana* was evaluated, demonstrating a significant delay in regeneration with reduced formation of new segments (41). Growth inhibition was observed in marine algae when *Isochrysis galbana* was exposed to 100 or 500 mg/L of caffeine (2). Polychaetes represented by *Hediste diversicolor*, *D. neapolitana*, and *Arenicola marina*, when exposed to caffeine concentrations between 0.5 and 18.0 µg/EU for 28 days, showed 8.3%, 12.5%, and 22.2% of lethality, respectively (41,42).

In the presence of caffeine, aquatic organisms increase their metabolic activity and reduce their energy reserves, such as glycogen, which are used to combat oxidative stress and prevent cell damage (15, 34, 41, 42). In addition to stimulating oxidative stress, caffeine can trigger apoptosis, as evidenced by the increased expression of Bax and Bcl2 (1). Exposure to caffeine, associated with other emerging pollutants, also alters the expression of genes in zebrafish that are used as markers of exposure to pollution (58). However, several questions remain regarding the genomic responses of aquatic organisms exposed to caffeine. For example, studies on the effects of caffeine on the expression of genes that control and participate in the development of aquatic organisms, including bone and cartilage development, are lacking.

Our team has been studying the effects, in mammals, of caffeine consumption by mothers on the bones of their offspring for several years. These studies have shown that caffeine inhibits postnatal bone growth and alters *in vivo* and *in vitro* expression of several gene transcripts related to osteogenesis and chondrogenesis (48-52). It has also been

shown that caffeine consumption by the mother can have teratogenic effects in the offspring (12). However, the effects of caffeine on embryogenesis have been poorly studied in mammals (65). In rats, because the endochondral bone formation is intrauterine, studies aiming to evaluate the effect of caffeine on this process supply the drug to the mother so that it reaches the fetus transplacentally. However, the lack of control over the amount of caffeine that reaches the fetus is the main limitation in using mammals as models to study the effects of caffeine or any other drug on endochondral bone formation. In this context, zebrafish are a more suitable model for studying bone changes resulting from failures in the bone formation process during embryonic development.

Despite the Zebrafish being widely used as a model to study the genesis of several human diseases, including cardiovascular, neuromuscular, and bone diseases (9, 11, 13, 36, 40, 46) and also be used as a model for studying the action of various drugs, including pollutants such as caffeine (2, 37, 43, 54), this seems to be to the best of our knowledge the first study that evaluated the effects of caffeine on the expression of gene transcripts related to osteogenesis and chondrogenesis, during zebrafish embryonic development.

Material and Methods

The procedures described below were approved by the Animal Use Ethics Committee of *Universidade Federal de Minas Gerais* under protocol nº. 115/2016.

Zebrafish maintenance and egg collection

Adult Zebrafish AB (*Danio rerio*) were raised in rearing rack systems using a filter system and water recirculation. The animals received granulated feed three times a day (4, 32) and were ordered in a regime of 12 hours of light and 12 hours of dark (13) at 28 ± 1°C. The pairs were placed in breeding tanks overnight so that newly laid eggs could be collected the next morning. Unfertilized or dead eggs were removed. Embryo staging was performed as described by Kimmel et al. (30).

Caffeine exposure

Fertilized eggs (embryos/5hpf) were collected and randomly divided into five experimental groups (128 eggs/embryo per group). Group 1: control (without caffeine); Group 2: 0.25 mM caffeine; Group 3: 0.5 mM caffeine; Group 4: 1 mM caffeine; and Group 5: 2 mM caffeine. The embryos were placed in 6-well cell culture plates containing water from the aquarium where spawning occurred and placed in greenhouses at a temperature of 28°C for up to 72 h post-fertilization. In the exposed groups, caffeine (Sigma-Aldrich,

St. Louis, MO, USA) in different concentrations was added to the water (35, 47, 61), and the embryos were exposed by immersion at 6 hpf. The medium was changed once daily (29).

External morphological changes in developing zebrafish embryos after caffeine exposed were carefully examined under a stereomicroscope (Stemi 508, Zeiss) at 6, 12, 24, 30, 48, 54, and 72 hpf, and images were captured using a digital camera (Axiocam 105 color, Zeiss).

Evaluation of gene transcript expression by real-time RT-PCR

The larvae of the five groups, 72hpf, were submitted to RT-PCR in real time for the quantitative evaluation of the expression of Sox9a, runx2b, osteocalcin, osteopontin, collagen 2a1 and bmp2a through protocols previously established (31). Larvae were macerated in Trizol. Specific PCR primer sequences were obtained from GenBank.

Gapdh: *forward*: CGCTGGCATCTCCCTCAA, *reverse*: TCAGCAACACGATGGCTGTAG (10.1111/j.1745-7270.2007.00283.x.); EFLa1: *forward*: CTGGAGGCCAGCTCAAACAT, *reverse*: ATCAAGAA-GAGTAGTACCGCTAGCATTAC (10.1111/j.1745-7270.2007.00283.x.); collagen 2a1: *forward*: CCTCTGAAATCCAGCCATGT, *reverse*: GACTGCTGTG-GTTCCAGTCA (<https://doi.org/10.1016/j.gep.2015.07.004>); bmp2a: *forward*: AGTGTTAGGACGACGCAGC, *reverse*: CGGCCTCATTTGAGACCAC (doi:10.1016/s0378-1119(97)00292-8.); runx2b: *forward*: ATGGCCGAGAT-CATCGCCGATCAC, *reverse*: CGGGGCCACCTG-GTTCTTCATAACC (doi.org/10.3389/fendo.2017.00125); sox9a: *forward*: CGGTGAAGAACGGCCAGAGC, *reverse*: CTGTAGAGTCAGCAATGGGT (doi.org/10.1006/dbio.2000.0129); osteocalcin: *forward*: TGGCCTCTAT-CATCATGAGACAGA, *reverse*: CTCTCGAGCTGAAAT-GGAGTCA (doi:10.1016/j.jot.2015.07.002); osteopontin: *forward*: GCCACCTCTATTCTTCG, *reverse*: TGCCCTC-CAGTGTCATCT (doi: 10.1007/s00223-013-9817-4).

Subsequently, lysis and homogenization were performed for 5 min at room temperature to completely dissociate the nucleoprotein complexes. The lysate was transferred to a 1.5mL microtube, with the addition of 0.2mL of chloroform, being homogenized for 15 seconds and incubated for 3 minutes at room temperature, followed by centrifugation at 12,000g for 15 minutes at 4°C, aiming to separate into three phases, the colorless and superficial phase, which contains the RNA. The aqueous phase was transferred to a new tube, with the addition of 0.5mL of isopropyl alcohol and incubation for 15 minutes at room temperature, followed by centrifugation at 12,000g for 10 minutes at 4°C for RNA precipitation. The pellet was washed with 1mL of 75% ethanol, homogenized, and centrifuged at 7,500g for 5 minutes at 4°C. The RNA was solubilized in RNase-free water and immediately stored at -80°C. The quality of the RNA samples was analyzed using spectrophotometry.

Reverse transcription reactions were performed using a commercial Taqman reverse transcription reagent kit (Applied Biosystems, Foster City, CA, USA). PCR was performed using 2µg of cDNA, 600nM of each primer, and 17µL of Syber Green reagent (Applied Biosystems, Foster City, CA, USA) in a final volume of 25µL of reaction in a 7900HT Fast Real-Time PCR System. The optimal conditions for the real-time PCR of the amplified cDNAs were determined in preliminary assays. The results obtained for each group were quantitatively compared after normalization based on GAPDH and EFLa1 danio rerio expression. The analysis was performed in quadruplicate.

Statistical analysis

Statistical analyses were performed using GraphPad Prism 8.0.2 software to statistically compare the experimental values between the control and each treatment group. The study design was randomized. Means and standard deviations were determined. Averages were compared using the Student's t-test. Data were tested for normality and homoscedasticity. Differences were considered statistically significant at $P < 0.05$.

Results

Effect of caffeine on the external appearance of developing zebrafish embryos/larvae

As previously described, normal embryonic development occurred in the control group (Fig. 1). At 6 hpf, the yolk sac, blastoderm, and chorion were observed. At 12 hpf, the beginning of the division of the embryonic planes (displacement of the head from the tail) was observed, concomitant with the appearance of the yolk syncytial layer. At 24 hpf and 30 hpf, the optic placode, pericardial cavity, somites, notochords, and olfactory placodes were observed. At 48 hpf stage, the embryo begins the process of melanization, the optic placode gives rise to the eyes, and the heart is observed. At 54 hpf the embryos started to hatch. At 72 hpf, all animals hatched were active and maintained a constant swimming position with spontaneous movement.

There was no significant difference in embryonic development in the exposed group with 0.25 mM of caffeine compared to the control group. The formation and development of embryo body structures followed the same temporal dynamics observed in the control group (Fig. 1). Spontaneous movement was unaffected by caffeine.

Morphological deformities in the embryos were observed in the 0.5 mM caffeine group at 12 hpf, where only the yolk sac, blastoderm, and chorion were observed. There




Figure 1. Gross morphological changes after caffeine treatment. The embryos were exposed to 0.25, 0.5, 1, and 2 mM caffeine from 6 to 72 hours post fertilization (hpf): 1: yolk sac; 2: corion; 3: head region; 4: tail region; 5: yolk syncytial layer; 6: notochord; 7: eye; 8: heart; 9: fin; T: tail. * axial-tail curvature; white arrow: pericardial edema; and red arrow: yolk sac malformations (edema).

was a delay at the beginning of the division of the embryo planes, which was different from that observed in the group with 0.25 mM of caffeine. However, at 24 hpf and 30 hpf, the optic placode, pericardial cavity, somites, notochords, and olfactory placodes were observed. At 48 hpf and 54 hpf, the embryos began the melanization process, the optic placode gave rise to the eyes, and the heart was observed. Some embryos began to hatch, and some animals displayed significant axial tail curvature. At 72 hpf, all animals hatched, and larvae without deformation were active in the swimming position with spontaneous movement. However, larvae with an axial tail curvature remained in a side-to-side position with less movement.

In the 1 mM group, external morphological malformations were more evident. Embryonic development was similar to that observed in animals treated with 0.5 mM up to 48 hpf. At 54 hpf, the animals did not start hatching and

exhibited significant axial tail curvature, pericardial edema, and yolk sac malformations (edema). At 72 hpf, these malformations were observed in all animals, such as those that hatched and those that did not (Fig. 1).

In the 2mM group, external morphological malformations were more evident. Embryonic development was similar to that observed in animals treated with 1 mM. Significant axial tail curvature, pericardial edema, and yolk sac malformation (edema) were observed as early as 48 hpf. At 72 hpf, the animal did not hatch (Fig. 1).

Expression of transcription factors associated with osteogenesis and chondrogenesis

Embryos exposed with 0.25 mM caffeine showed significantly lower relative expression of *bmp2a*, *runx2b*, and

collagen 2a1 compared to the control group; however, there was no significant difference in the expression of osteocalcin, osteopontin, and Sox9a (Fig. 2). In the 0.5 mM caffeine group, the relative expression levels of *bmp2a*, *runx2b*, and collagen 2a1 were significantly lower than those in the control, while osteocalcin expression was significantly higher. No significant differences were observed in the expression of osteopontin and *sox9* (Fig. 2). In the 1 mM caffeine group, the relative expression levels of *bmp2a*, *runx2b*, and collagen 2a1 were significantly lower than those in the control group, and the osteocalcin expression was significantly higher. No significant differences were observed in the expression of osteopontin and *sox9a* (Fig. 2). In the 2mM caffeine group, the relative expression of *bmp2* and *runx2* was significantly lower, whereas that of osteocalcin, osteopontin, *sox9a*, and collagen 2a1 did not differ from that of the control group (Fig. 2).

Discussion

Based on the results presented, the genomic effects of caffeine on the expression of gene transcripts related to chondrogenesis and osteogenesis can occur independently of embryonic morphological changes; that is, the addition of caffeine to the medium containing zebrafish embryos can alter the expression of gene transcripts, even without altering embryonic morphology.

This result was visible at 0.25 mM caffeine, which significantly reduced the expression of *bmp2a*, *runx2b*, and collagen 2a1, without altering the embryonic morphology, as observed at higher concentrations of caffeine. Caffeine

significantly reduced the expression of *bmp2a* and *runx2b* transcripts at all concentrations. At concentrations lower than 2 mM, there was also a reduction in collagen 2a1. The transcriptional alteration of genes involved in chondrogenesis and osteogenesis promotes a better understanding of the mechanisms involved in changes in bone development caused by caffeine.

The caffeine concentrations were determined based on previous studies that tested similar concentrations in zebrafish (16, 22, 45, 60) and rat chondrocytes. In addition to changes in the notochord and tail, the embryonic morphological changes found at concentrations of 0.5 mM, 1 mM, and 2 mM caffeine were not directly related to the process of chondrogenesis and osteogenesis but reaffirmed the toxic and teratogenic potential of caffeine (9, 45, 46, 59). Alterations in gene transcripts related to caffeine-induced chondrogenesis and osteogenesis during embryonic development have not been previously described. One of the interesting results of this study is the evidence that caffeine at a concentration of 0.25 mM can alter the profile of these gene transcripts in apparently healthy embryos.

Caffeine-induced changes in the expression of gene transcripts without phenotypic changes are interesting because some studies have shown that zebrafish embryos descending from parental exposure to environmental pollutants may exhibit developmental abnormalities, even if they did not occur in the parents. These subsequent generational effects, mediated by mothers and fathers exposed to toxicants, may occur due to the bioaccumulation and transfer of toxicants in adult zebrafish germ cells. However, there is evidence that developmental changes occur through epigenetic modification




Figure 2. Expression of gene transcripts (mean \pm standard deviation) for *bmp2a*, *runx2b*, osteopontin, osteocalcin, *sox9a*, and Col21a by the RT-qPCR in zebrafish embryos treated with 0.25; 0.5; 1, and 2 mM caffeine at 72hpf. $p \leq 0.05^*$

of genes transferred from F0 to F1 larvae (67). Thus, complementary transgenerational studies are required to verify whether the transcriptional changes induced by caffeine can be passed on to subsequent generations.

The effect of caffeine on gene expression has been studied in the embryo, as chondrogenesis and osteogenesis begin in the embryonic phase of mammals and teleosts. In zebrafish, this process begins during embryonic development before hatching. The zebrafish skeleton has a simple organization composed of cartilage and bone, with the craniofacial cartilage being one of the first to form during development. The first elements, represented by the branchial arch and opercular bones, develop 48 h after fertilization, generating a fully functional cartilaginous skeleton on the fifth day after fertilization, that is, two days after hatching of the larva (23, 30, 55, 64). Research on gene deletion in zebrafish has shown that several genes, including those studied here, such as *runx2ab*, *sox9a*, *collagen 2a1*, and *bmp2a*, also participate in chondroblast differentiation in the craniofacial region of zebrafish, similar to that in mammals (10, 24, 44, 63).

All caffeine concentrations used significantly reduced gene transcript expression for *runx2b*. *Runx2* is an essential transcription factor for osteoblast, chondrocyte differentiation, and cartilage and bone development in vertebrates (17). In mammals, *runx2* has two distinct functions: one is essential for the differentiation of mesenchymal progenitor cells into osteoblasts, and the other stimulates the differentiation of hypertrophic chondrocytes (14) and is therefore considered a key regulator of osteogenesis (8).

In zebrafish, the *runx* genes exhibit a hierarchy of functions during skeletogenesis. *runx2b* plays a dominant role in differentiating pre-cartilaginous cells into chondrocytes. Cephalic endochondral osteogenesis begins 48 h after fertilization via focal mesenchymal condensation (55). At this stage, both *runx2a* and *runx2b* are expressed in the primordia of dermal and cartilaginous bones. However, unlike mammals, *runx1* is not important in the early stages of chondrocyte differentiation. In contrast, *runx2b* depletion seriously compromises the formation of the craniofacial ethmoid cartilage (24). Therefore, it is possible that the reduction in expression of the gene transcript for *runx2b* caused by all caffeine concentrations subsequently determines some type of alteration in the bones that develop from the structures in which it is expressed.

Caffeine effects on *runx2* expression in mice or rat differ from those in zebrafish. Osteoblasts extracted from the calvariae of rats born to mothers who received caffeine during pregnancy significantly increased *runx2* expression. Increased *runx2* expression is associated with increased osteoblastic activity and the expression of other gene transcripts such as osteocalcin, osteopontin, sialoprotein, and collagen 1 (50). However, the expression of *runx2* in chondrocytes extracted from rats born to mothers who received caffeine during pregnancy did not differ from that in the untreated group (control) (52). The effects of caffeine on mammalian and zebrafish

expression of *runx 2* differ, possibly due to several reasons besides the inherent differences in animal species. In rats, the effect of caffeine has been studied after birth (50, 52) but not during embryonic development, as in the present study. Furthermore, in studies by Reis et al. (50, 52), caffeine was transferred to the fetus through the placenta, and the effect of caffeine was studied on a specific population of cells, in this case, osteoblasts, and chondrocytes.

In addition to reducing *runx2b* expression, all the caffeine concentrations significantly reduced gene transcript expression for *bmp2a*. *Bmp2* plays an important role in regulating osteoblast proliferation and differentiation. In zebrafish, *bmp2a* and *bmp2b* are primarily expressed in calcified tissues (38). Knockdown of the antisense morpholino oligonucleotide gene mediated by *bmp2a* causes a delay in early bone development and decreases the expression of osteogenic genes, such as *runx2a* (28). Mutations in *bmp2a* in zebrafish compromise complete rib formation and cause vertebral deformities (66). It is suggested that the reduced expression of the *bmp2a* transcript may be one of the reasons why zebrafish larvae show defects, such as tail curling. However, this hypothesis warrants further investigation.

At caffeine concentrations lower than 2 mM, there was a reduction in the transcript of the gene encoding collagen 2a1. Collagen is the most abundant constituent of the chondrogenic matrix. During chondrogenesis in mammals, the expression of different types of collagens occurs temporally according to the level of cell differentiation, with the initial expression of collagen type 1 by undifferentiated cells, followed by collagen type 2 by differentiated cells, and finally, by collagen type 10 in mature cells (5, 56, 57). In zebrafish, no studies have demonstrated the temporal profile of collagen expression during development. However, 2a1 collagen gene expression has been identified in the craniofacial cartilage, ear, notochord, floor plate, hypochondrium, and zebrafish fin. In this fish, 2a1 collagen expression is a marker of cartilage and notochord development (18). It is likely that, in addition to the reduction of the gene transcripts for *runx2b* and *bmp2a*, the reduction of the transcript for collagen 2a1 is also involved in the genesis of the alterations caused by caffeine, not only in the skeletal system but also in other tissues and organs containing this type of collagen.

None of the caffeine concentrations studied altered the gene transcript expression for *sox9a*. *Sox9* stimulates the differentiation of chondroprogenitor cells into chondroblasts (19, 33). The inactivation of *sox9* in mice results in the absence of expression of chondrogenic markers, such as collagens 2, 9, and 11, and aggrecan, which prevents cartilage formation (7). Like mammals, *sox9a* and *sox9b* are regulators of cartilage development and are necessary for the expression of collagen 2a1; therefore, they are essential for endochondral bone formation in zebrafish (17). Based on the results of the present study, it can be concluded that at the concentrations studied and up to 78 h after fertilization, caffeine did not alter the expression of *sox9a*.

However, additional studies are required to verify whether caffeine alters the expression of this transcript in the later stages of post-embryonic development. In mammals, we found antagonistic effects of caffeine on *sox9* expression in chondrocytes. The addition of caffeine to a culture medium with cartilaginous epiphyses of the femurs of newborn rats significantly increased the expression of the transcript for *sox9* in chondrocytes (48). In contrast, the articular cartilage of rats born to mothers who received caffeine during pregnancy significantly reduced *sox9* expression (52).

None of the caffeine concentrations significantly altered osteopontin expression. However, treatment with 1 mM caffeine significantly increased osteocalcin expression. In mammals, osteopontin is expressed during cell proliferation and at the beginning of mineralization (68), promoting the premature precipitation of calcium phosphate crystals in the collagen matrix (20). Osteocalcin, on the other hand, is found later in the course of the bone mineralization process (3, 53). In zebrafish, although osteocalcin is expressed in the vertebral bodies, one of the osteocalcin isoforms, *oc1*, is expressed in the notochord before osteoblastic differentiation and bone formation. Thus, it is likely that *oc1* is involved in processes before bone formation, that is, in the mineralization of the notochord sheath.

The notochord is important in the early stages of life because its mechanical function is only replaced by the vertebral column in the post-embryonic phase (6). Based on this study, it is difficult to understand why only 1 mM caffeine significantly increased osteocalcin expression, despite significantly reducing the expression of transcripts for *bmp2*, *runx2*, and collagen 2. The implications of this caffeine-induced increase in osteocalcin expression on notochord mineralization and subsequent skeletal development require further study. Osteoblasts extracted from the calvariae of rats born to mothers who received caffeine during pregnancy showed a significant increase in osteocalcin expression. However, contrary to the results, increased osteocalcin expression was associated with increased expression of other gene transcripts, such as osteopontin, sialoprotein *runx-2*, and collagen 1 (50).

Conflict of Interest

The authors declare no competing interests.

Acknowledgments

This work was supported by grants from the Fundação de Amparo à Pesquisa de Minas Gerais (Fapemig: APQ 01115-17), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

References

1. Abdelkader TS, Chang SN, Kim TH, Song J, Kim DS, Park JH. Exposure time to caffeine affects heartbeat and cell damage-related gene expression of zebrafish *Danio rerio* embryos at early developmental stages. *J Appl Toxicol*. 2013; 33(11):1277–1283.
2. Aguirre-Martínez GV, Reinardy HC, Martín-Díaz ML, Henry TB. Response of gene expression in zebrafish exposed to pharmaceutical mixtures: Implications for environmental risk. *Ecotoxicol Environ Saf*. 2017; 142:471–479.
3. Aksu AE, Rubin JP, Dudas JR, Marra KG. Role of gender and anatomical region on induction of osteogenic differentiation of human adipose-derived stem cells. *Ann Plast Surg*. 2008; 60(3):306–322.
4. Avdesh A, Chen M, Martin-Iverson MT, Monda LA, Ong D, Rainey-Smith S, Taddei K, Lardelli M, Groth DM, Verdile J, Martins RN. Regular care and maintenance of a zebrafish (*Danio rerio*) laboratory: An introduction. *J Visual Exp*. 2012; 18(69):1–8.
5. Barry F, Boynton RE, Liu B, Murphy JM. Chondrogenic differentiation of mesenchymal stem cells from bone marrow: differentiation-dependent gene expression of matrix components. *Exp Cell Res*. 2001; 268(2):189–200.
6. Bensimon-Brito A, Cancela ML, Huisseune A, Witten PE. Vestiges, rudiments, and fusion events: the zebrafish caudal fin endoskeleton in an evo-devo perspective. *Evol Dev*. 2012; 14(1):116–127.
7. Bi W, Deng JM, Zhang Z, Beringer RB, Crombrughe B. *Sox9* is required for cartilage formation. *Nat Genet*. 1999; 22(1):85–89.
8. Bruderer M, Richards RG, Alini M, Stoddart MJ. Role and regulation of *RUNX2* in osteogenesis. *Eur Cell Mater*. 2014; 28:269–286.
9. Chiang EF, Pai CI, Wyatt M, Yan YL, Postlethwait J, Chung B. Two *sox9* genes on duplicated zebrafish chromosomes: expression of similar transcription activators in distinct sites. *Dev Biol*. 2001; 231(1):149–63.
10. Chen YH, Huang, YH, Wen CC, Wang YH, Chen WL, Chen LC, Tsay H. Movement disorder and neuromuscular change in zebrafish embryos after exposure to caffeine. *Neurotoxicol Teratol*. 2008; 30(5):440–7.
11. Chico TJ, Ingham PW, Crossman DC. Modeling cardiovascular disease in the zebrafish. *Trends Cardiovasc Med*. 2008; 18(4):150–5.
12. Christian MS, Brent RL. Teratogen update: evaluation of the reproductive and developmental risks of caffeine. *Teratol*. 2001; 64(1):51–78.
13. Cooper WJ, Wirgau RM, Sweet EM, Albertson RC. Deficiency of zebrafish *fgf20a* results in aberrant skull remodeling those mimics both human cranial disease and evolutionarily important fish skull morphologies. *Evol Dev*. 2013; 15(6):426–441.

14. Cruz D, Almeida A, Calisto V, Esteves VI, Schneider RJ, Wrona FJ, Soares AM, Figueira E, Freitas R. Caffeine impacts in the clam *Ruditapes philippinarum*: alterations on energy reserves, metabolic activity, and oxidative stress biomarkers. *Chemosphere* 2016; 160: 95e103.
15. Crombrughe B, Lefebvre V, Nakashima K. Regulatory mechanisms in the pathways of cartilage and bone formation. *Curr Opin Cell Biol.* 2001; 13(6):721-727.
16. Cruz FF, Leite C, Kist LW, Oliveira GM, Bogo MR, Bonan C, Campos M, Morrone FB. Effects of caffeine on behavioral and inflammatory changes elicited by copper in zebrafish larvae: Role of adenosine receptors. *Comp Biochem Physiol. Part C* 2017; 194:28-36.
17. Dalcq J, Pasque V, Ghaye A, Larbuisson A, Motte P, Martial JA, Muller M. RUNX3, EGR1 and SOX9B Form a Regulatory Cascade Required to Modulate BMP-Signaling during Cranial Cartilage Development in Zebrafish. *PLoS One* 2012; 7(11):e50140.
18. Dale RM, Topczewski J. Identification of an evolutionarily conserved regulatory element of the zebrafish *col2a1a* gene. *Dev Biol.* 2011; 357(2): 518-531.
19. Delise AM, Fischer L, Tuan RS. Cellular interactions and signaling in cartilage development. *Osteoarthritis Cartil.* 2000; 8(5):309-334.
20. Doi Y, Horiguchi T, Kim SH, Moriwaki Y, Wakamatsu N, Adachi M, Ibaraki K, Moriyama K, Sasaki S, Shimokawa H. Effects of non-collagenous proteins of the formation of apatite in Ca- β -glycerophosphate solution. *Arch Oral Biol.* 1992; 37(1):15-21.
21. Dubey M, Vellanki BP, Kazmi AA. Removal of emerging contaminants in conventional and advanced biological wastewater treatment plants in India-a comparison of treatment technologies. *Environ Res.* 2023; 218:115012.
22. Felix L, Lobato-Freitas C, Monteiro SM, Venancio C. 24-Epibrassinolide modulates the neurodevelopmental outcomes of high caffeine exposure in zebrafish (*Danio rerio*) embryos. *Comp Biochem Physiol Part C* 2021; 249:109143.
23. Fiaz AW, Leon-Kloosterziel KM, Gort G, Schulte-Merker S, vanLeeuwen JL, Kranenbarg S. Swim-Training Changes the Spatio-Temporal Dynamics of Skeletogenesis in Zebrafish Larvae (*Danio rerio*). *Plos one* 2012; 7(4):e34072.
24. Flores MV, Lam EY, Crosier P, Crosier K. A hierarchy of Runx transcription factor modulates the onset of chondrogenesis in craniofacial endochondral bones in zebrafish. *Dev Dyn.* 2006; 235(11):3166-76.
25. French VA, King SC, Kumar A, Northcott G, McGuinness K, Parry D. Characterisation of microcontaminants in Darwin Harbour, a tropical estuary of northern Australia undergoing rapid development. *Sci Total Environ.* 2015; 536:639-647.
26. Garcia RN, Chung KW, Delorenzo ME, Curran MC. Individual and mixture effects of caffeine and sulfamethoxazole on the daggerblade grass shrimp *Palaemonetes pugio* following maternal exposure. *Environ Toxicol Chem.* 2014; 33(9):2120-2125.
27. Gonzalez-Rey M, Tapie N, Le Menach K, Devier MH, Budzinski H, Bebianno MJ. Occurrence of pharmaceutical compounds and pesticides in aquatic systems. *Mar Pollut Bull.* 2015; 96(1-2):384-400.
28. Jiang Y, Wang L, Zhu GX, Xu YJ. Effects on osteoblasts by bone morphogenetic protein 2a knocked down in zebrafish. *CJOBMR* 2019; 12:165-171.
29. Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. Stages of Embryonic Development of the Zebrafish. *Dev Dyn.* 1995; 203(3):253-310.
30. Kimmel CB, Miller CT, Moens CB. Specification and morphogenesis of the zebrafish larval head skeleton. *Dev Biol.* 2001; 233(2):239-257.
31. Lan CC, Tang R, Leong IUS, Love DR. Quantitative real-time RT-PCR (qRT-PCR) of zebrafish transcripts: optimization of RNA extraction, quality control considerations, and data analysis. *Cold Spring Harb Protoc.* 2009; 10: 5314.
32. Lawrence C. The husbandry of zebrafish (*Danio rerio*): A review. *Aquac.* 2007; 269(1-4):1-20.
33. Lefebvre V, Li P, Crombrughe B. A new long form of Sox-5 (L-Sox-5), Sox-6 and Sox-9 are coexpressed in chondrogenesis and cooperatively activate the type II collagen gene. *Embo J.* 1998; 17(19):5718-5733.
34. Li S, He B, Wang J, Liu J, Hu X. Risks of caffeine residues in the environment: Necessity for a targeted ecopharmacovigilance program. *Chemosphere*, 2020; 243:125343.
35. Lu PZ, Lai CY, Chan WH. Caffeine induces cell death via activation of apoptotic signal and inactivation of survival signal in human osteoblasts. *Int J Mol Sci.* 2008; 9(5):698-718.
36. Marí-Beffa M, Santamaría JA, Murcian C, Santos-Ruiz L, Andrade JA, Guerado E, Becerra J. Zebrafish Fins as a Model System for Skeletal Human Studies. *Sci World J.* 2007; 7:1114-1127.
37. Martini GA, Montagner CC, Viveiros W, Quinaglia GA, França DD, Munin NCG, Lopes-Ferreira M, Rogero SO, Rogero JR. Emerging contaminant occurrence and toxic effects on zebrafish embryos to assess the adverse effects caused by mixtures of substances in the environment. *Environ Sci Pollut Res.* 2021; 28(16):20313-20329.
38. Marques CL, Fernandez I, Viegas MN, Cox CJ, Martel P, Rosa J, Cancela ML, Laiz EV. Comparative analysis of zebrafish bone morphogenetic proteins 2, 4 and 16: molecular and evolutionary perspectives. *Cell Mol Life Sci.* 2016; 73(4):841-857.
39. Nodler K, Voutsas D, Licha T. Polar organic micropollutants in the coastal environment of different marine systems. *Mar Pollut Bull.* 2014; 85(1):50-59.

40. Pappalardo A, Pitto L, Fiorillo C, Donati M, Bruno C, Santorelli FM. Neuromuscular disorders in zebrafish: state of the art and future perspectives. *Neuromolecular Med.* 2013; 15(2):405-19.
41. Pires A, Almeida A, Calisto V, Schneider RJ, Esteves VI, Wrona FJ, Soares AMVM, Figueira E, Freitas R. Long-term exposure of poly- chaetes to caffeine: biochemical alterations induced in *Diopatra neapolitana* and *Arenicola marina*. *Environ Pollut.* 2016a; 214:456-463.
42. Pires A, Almeida A, Correia J, Calisto V, Schneider RJ, Esteves VI, Soares AM, Figueira E, Freitas R. Long-term exposure to caffeine and carbamazepine: impacts on the regenerative capacity of the polychaete *diopatra neapolitana*. *Chemosphere.* 2016b; 146:565-573.
43. Pruvot B, Quiroz Y, Voncken A, Jeanray N, Piot A, Martial JA, Muller M. A panel of biological tests reveals developmental effects of pharmaceutical pollutants on late stage zebrafish embryos. *Reprod Toxicol.* 2012; 34(4):568-583.
44. Quint E, Smith A, Avaron F, Laforest L, Miles J, Gaffield W, Akimenko M. Bone patterning is altered in the regenerating zebrafish caudal fin after ectopic expression of sonic hedgehog and *bmp2* or exposure to cyclopamine. *Proc Natl Acad Sci USA.* 2002; 99(13):8713-8.
45. Rah YC, Yoo MH, Choi J, Park S, Park HC, Oh KH, Lee SH, Kwon SY. In vivo assessment of hair cell damage and developmental toxicity caused by gestational caffeine exposure using zebrafish (*Danio rerio*) models. *Neurotoxicol Teratol.* 2017; 64:1-7.
46. Rana N, Moond M, Marthi A, Bapatla S, Sarvepalli T, Chatti K, Challa AK. Caffeine-induced effects on heart rate in zebrafish embryos and possible mechanisms of action: an effective system for experiments in chemical biology. *Zebrafish.* 2010; 7(1):69-81.
47. Rapuri PB, Gallagher JC, Nawaz Z. Caffeine decreases vitamin D receptor protein expression and 1,25(OH)₂D₃ stimulated alkaline phosphatase activity in human osteoblast cells. *J Steroid Biochem Mol Biol.* 2007; 103(3-5):368-371.
48. Reis AMS, Raad RV, Ocarino NM, Serakides R. Efeitos in vitro da cafeína na cartilagem de crescimento de ratos. *Acta Ortop Bras.* 2013; 21(6):307-309.
49. Reis AMS, Batista ACM, Ocarino NM, Serakides R. Effects of caffeine intake in mothers on maternal cortisol levels and offspring endochondral ossification. *Atualizações em osteologia.* 2014; 10(1):20-36.
50. Reis AMS, Ribeiro LGR, Ocarino NM, Serakides R. Osteogenic potential of osteoblasts from neonatal rats born to mothers treated with caffeine throughout pregnancy. *BMC Musculoskelet Disord.* 2015; 16(1):1-11.
51. Reis AMS, Ocarino NM, Boeloni JN, Gomes DA, Goes AM, Ferreira AF, Serakides R. Inhibition of the osteogenic differentiation of mesenchymal stem cells derived from the offspring of rats treated with caffeine during pregnancy and lactation. *Connect Tissue Res.* 2016; 57(2):1-12.
52. Reis AMS, Oliveira KP, Paula IHF, Silva AP, Tarragô JF, Ocarino NM, Serakides R. Nonlinear effects of caffeine on the viability, synthesis, and gene expression of chondrocytes from the offspring of rats treated during pregnancy. *Acta Histochem.* 2018; 120(6):505-512.
53. Roach HI. Why does bone matrix contain non-collegeneous protein? The possible roles of osteocalcin, osteonectin, osteopontin and bone sialoprotein in bone mineralization and reabsorption. *Cell Biol Int.* 1994; 18(6):617-628.
54. Santos N, Picolo V, Domingues I, Perillo V, Villacis RAR, Grisolia CK, Oliveira M. Effects of environmental concentrations of caffeine on adult zebrafish behaviour: a short-term exposure scenario. *Environ Sci Pollut Res.* 2023; 30(23):63776-63787.
55. Schilling TF, Kimmel CB. Musculoskeletal patterning in the pharyngeal segments of the zebrafish embryo. *Dev.* 1997; 124(15):2945-60.
56. Shum L, Coleman CM, Hatakeyama Y, Tuam R. Morphogenesis and dysmorphogenesis of the appendicular skeleton. *Birth Defects Res Part C: Embryo Today.* 2003; 69(2):102-122.
57. Singh P, Schwarzbauer JE. Fibronectin and stem cell differentiation - lessons from chondrogenesis. *J Cell Sci.* 2012; 125(Pt16):3703-12.
58. Sposito JCV, Montagner CC, Casado M, Navarro-Martín L, Solórzano JCJ, Piña B, Grisolia AB. Emerging contaminants in Brazilian rivers: Occurrence and effects on gene expression in zebrafish (*Danio rerio*) embryos. *Chemosphere.* 2018; 209:696-704.
59. Teixido E, Pique E, Gomez-Catalan J, Llobet JM. Assessment of developmental delay in the zebrafish embryo teratogenicity assay. *Toxicol In Vitro.* 2013; 27(1):469-478.
60. Tran S, Fulcher N, Nowicki M, Desai P, Tsang B, Faciol A, Chow H, Gerali R. Time-dependent interacting effects of caffeine, diazepam, and ethanol on zebrafish behaviour. *Prog Neuropsychopharmacol Biol Psychiatry.* 2017; 75:16-27.
61. Tsuang JSS, Chen LT, Sun SCK, Chen SC. Direct effects of caffeine on osteoblastic cells metabolism: the possible causal effect of caffeine on the formation of osteoporosis. *J Orth Surg Res.* 2006; 1(7):1-7.
62. Vieira LR, Soares AMVM, Freitas R. Caffeine as a contaminant of concern: A review on concentrations and impacts in marine coastal systems. *Chemosphere* 2022; 286(Pt2):131675.
63. Yan YL, Miller CT, Nissen RM, Singer A, Liu D, Kirn A, Draper B, Willoughby J, Morcos PA, Amesdam A, Chung B, Westerfield M, Haffter P, Hopkins N, Kimmel C, Postlethwait JH. A zebrafish *sox9* gene required for cartilage morphogenesis. *Dev.* 2002; 129(21):5065-79.
64. Yelick PC, Schilling TF. Molecular dissection of craniofacial development using zebrafish. *Crit Rev Oral Biol Med.* 2002; 13(4):308-22.

65. Zhang Y, Chen Y, Xu K, Fang L, Huang J, Xia S, Zhou Q, Lv L, Wang C. Embryonic exposure to phenanthrene caused developmental defects of craniofacial cartilage in F1 larvae. *Aquat Toxicol.* 2022; 243:106080
66. Wilkinson JM, Pollard I. In utero exposure to caffeine causes delayed neural tube closure in rat embryo. *Teratog Carcinog Mutagen.* 1994; 14(5):205–211.
67. Wu Y, Sun A, Nie C, Gao Z, Wan S. Functional differentiation of *bmp2a* and *bmp2b* genes in zebrafish. *Gene Expression Patterns* 2022; 46:119288.
68. Zohar R, Cheifetz S, Mcculloch CA, Sodek J. Analysis of intracellular osteopontin as a marker of osteoblastic cell differentiation and mesenchymal cell migration. *Eur J Oral Sci.* 1998; 106(suppl1):401-407.